



## Product Datasheet

<b>Product Name</b>	Regenerating Protein 1 alpha Human Recombinant
<b>Cata No</b>	CB500910
<b>Source</b>	<i>Escherichia Coli.</i>
<b>Synonyms</b>	Lithostathine-1-alpha, Pancreatic stone protein, PSP, Pancreatic thread protein, PTP, Islet of Langerhans regenerating protein, REG, Regenerating protein I alpha, Islet cells regeneration factor, ICRF, REG1A, PSPS, P19, PSPS1, MGC12447.

### Description

REG protein was shown to be stimulated during the regeneration of pancreatic islets. Since then, many Reg-related proteins have been identified in humans and other animals. In human, the four REG family genes, i.e., REG 1 alpha, REG 1 beta, REG-related sequence (RS) and HIP/PAP, have so far been isolated. These Reg-related proteins are classified into four subfamilies according to their amino-acid sequences, but they share a similar structure and physiological function. Reg protein is a growth factor for pancreatic beta cells and also suggests that the administration of Reg protein could be used as another therapeutic approach for diabetes mellitus. Human REG cDNA which encodes a 166-amino acid protein with a 22-amino acid signal peptide. The amino acid sequence of human REG protein has 68% homology to that of rat Reg protein. Reg I was found to be expressed mainly in pancreatic beta and acinoductular cells as well as gastric fundic enterochromaffin-like (ECL) cells. Reg I production in ECL cells is stimulated by gastrin, as well as by the proinflammatory cytokine, cytokine-induced neutrophil chemoattractant (CINC)-2Beta. In patients with chronic hypergastrinemia, Reg production is stimulated, with the increased proliferation of gastric mucosal cells. Patients with Helicobacter pylori infection also showed increased Reg production in the gastric mucosa, partly via increased plasma gastrin

concentration and partly via increased proinflammatory cytokine production. The serum concentration of the reg-protein was significantly higher in patients with various pancreatic diseases than in normal controls, and was also significantly higher in patients with acute pancreatitis or chronic relapsing pancreatitis than in patients with chronic pancreatitis. Furthermore, the serum PSP/reg-protein concentration was also significantly increased in liver cirrhosis, choledocholithiasis, and various cancers of the digestive system.

The Recombinant Human REG 1 alpha is produced with N-terminal fusion His Tag. The Recombinant Human REG 1 alpha His-Tagged Fusion Protein, has a molecular weight of 17.8 kDa protein containing 144 amino acid residues of the Human REG 1 alpha and 12 additional amino acid residues – His Tag (underlined).

MKHHHHHHAS HMQEAQTELP QARISCEP  
GT NAYRSYCYFF NEDRETWVDA DLYCQNMNSG  
NLVSVLTQAE GAFVASLIKE SGTDDFNVWI  
GLHDPKKNRR WHWSSGSLVS YKSWGIGAPS  
SVNPGYCVSL TSSTGFQKWK DVPCEDKFSF  
VCKFKN

### Purity

Greater than 95% as determined by SDS-PAGE.

### Formulation

Sterile filtered and lyophilized from 0.5 mg/ml in 5mM Tris, 25mM NaCl, pH 7.5.

**\* For Non-Clinical Research Use Only \***



# California Bioscience

83103 Avenue 48, Ste.1B #204  
Coachella, CA 92236 USA  
Phone : +1.6268339877  
Email : [info@cali-bio.com](mailto:info@cali-bio.com)

## Reconstitution

Add 0.2 ml of deionized H<sub>2</sub>O and let the lyophilized pellet dissolve completely.

## Stability

Store lyophilized protein at -20°C. Aliquot the product after reconstitution to **avoid repeated freezing/thawing cycles**. Reconstituted protein can be stored at 4°C for a limited period of time; it does not show any change after two weeks at 4°C.

The lyophilized protein remains stable until the expiry date when stored at -20°C.

## Product Data sheet

## Applications

Western blotting, ELISA

**\* For Non-Clinical Research Use Only \***